

Assessment of Habitat of Wildlife Communities on the Snake River, Jackson, Wyoming



UNITED STATES DEPARTMENT OF THE INTERIOR
Fish and Wildlife Service / *Resource Publication 190*

DTIC 970319 040

Technical Report Series

U.S. Fish and Wildlife Service

The Fish and Wildlife Service publishes five technical report series. Manuscripts are accepted from Service employees or contractors, students and faculty associated with cooperative fish and wildlife research units, and other persons whose work is sponsored by the Service. Manuscripts are received with the understanding that they are unpublished. Most manuscripts receive anonymous peer review. The final decision to publish lies with the editor.

Series Descriptions

Editorial Staff

MANAGING EDITOR

Paul A. Opler

ASSISTANT MANAGING EDITOR

Paul A. Vohs

WILDLIFE EDITOR

Elizabeth D. Rockwell

FISHERIES EDITOR

James R. Zuboy

PUBLICATIONS MANAGEMENT

Thomas J. Cortese

TECHNICAL EDITORS

Deborah K. Harris, Senior Editor

John S. Ramsey

VISUAL INFORMATION SPECIALIST

Constance M. Lemos

EDITORIAL ASSISTANTS

Rhonda G. Brown

Amy D. Trujillo

North American Fauna ISSN 0078-1304

Monographs of long-term or basic research on faunal and floral life histories, distributions, population dynamics, taxonomy, and community ecology.

Fish and Wildlife Research ISSN 1040-2411

Scientific papers of scholarly quality on original research, theoretical presentations, and interpretive literature reviews.

Biological Report ISSN 0895-1926

Technical papers about applied research of a more limited scope than papers in *Fish and Wildlife Research*. Subjects include new information arising from comprehensive studies, surveys and inventories, effects of land use on fish and wildlife, diseases of fish and wildlife, animal control, and developments in technology. Proceedings of technical conferences and symposia may be published in this series.

Resource Publication ISSN 0163-4801

Semitechnical and nonexperimental technical topics including surveys; data, status, and historical reports; handbooks; checklists; manuals; annotated bibliographies; and workshop papers.

Fish and Wildlife Leaflet ISSN 0899-451X

Summaries of technical information for readers of nontechnical or semitechnical materials. Subjects include topics of current interest, results of inventories and surveys, management techniques, and descriptions of imported fish and wildlife and their diseases.

Copies of this publication may be obtained from the Publications Unit, U.S. Fish and Wildlife Service, 1849 C Street, N.W., Arlington Square Building, Mail Stop 130, Washington, DC 20240, or may be purchased from the National Technical Information Service (NTIS), 5285 Port Royal Road, Springfield, VA 22161 (Call Toll Free 1-800-553-6847).

Assessment of Habitat of Wildlife Communities on the Snake River, Jackson, Wyoming

By Richard L. Schroeder
Arthur W. Allen

UNITED STATES DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE
Resource Publication 190
Washington, D.C. • 1992

DRUG QUALITY INSPECTED 1

Contents

	Page
Abstract	1
Characteristics of Riparian Wildlife Habitat	2
Factors of Wildlife Diversity in Riparian Habitats	3
Structure and Abundance of Riparian Vegetation	3
Channel and Floodplain Complexity	5
Spatial Relations	8
Summary of Riparian Habitat and Wildlife Diversity	8
Habitat Suitability Index Model for a Riparian Wildlife Community	9
Model Overview and Output	9
Model Applicability	9
Model Variables and Assumptions	9
Habitat Suitability Index Determination and Application of the	
Model	13
References	15
Appendix. Riparian-dependent Vertebrates from the Region of the	
Snake River near Jackson, Wyoming	20

Assessment of Habitat of Wildlife Communities on the Snake River, Jackson, Wyoming

by

Richard L. Schroeder and Arthur W. Allen

*U.S. Fish and Wildlife Service
National Ecology Research Center
4512 McMurtry Avenue
Fort Collins, Colorado 80525*

Abstract. The composition of the wildlife community in western riparian habitats is influenced by the horizontal and vertical distribution of vegetation, the physical complexity of the channel, and barriers to movement along the corridor. Based on information from the literature and a workshop, a model was developed to evaluate the wildlife community along the Snake River near Jackson, Wyoming. The model compares conditions of the current or future years with conditions in 1956, before the construction of levees along the river. Conditions in 1956 are assumed to approximate the desirable distribution of plant cover types and the associated wildlife community and are used as a standard of comparison in the model. The model may be applied with remotely sensed data and is compatible with a geographic information systems analysis. In addition to comparing existing or future conditions with conditions in 1956, the model evaluates floodplain and channel complexity and assesses anthropogenic disturbance and its potential effect on the quality of wildlife habitat and movement of wildlife in the riparian corridor.

Key words: Riparian, community, wildlife, habitat, model.

This document provides a method and useful information for assessing changes and mitigating adverse changes of wildlife habitat along 40 km of the Snake River near Jackson, Wyoming. The changes were created by the Jackson Hole-Snake River levee system that was constructed between the early 1950's and the late 1970's.

On 25 and 26 June 1991, a workshop was held to develop an assessment technique for evaluating changes in the wildlife community of the affected reaches of the Snake River near Jackson, Wyoming. The workshop concept was proposed by the tri-agency team (U.S. Fish and Wildlife Service

[Service], Wyoming Game and Fish Department, and U.S. Army Corps of Engineers) responsible for evaluating the effects of the levees and was supported by the National Ecology Research Center of the Service. The objective of the workshop was to develop a community-level, quantitative model for evaluating wildlife habitat in the floodplain with primarily remotely sensed data. The desire for a community-level model stems from an interest in moving beyond assessment and management of habitat for single species. As noted by Schroeder (1987), the interest in community-level approaches for wildlife-habitat assessment is growing. To be of

most use, assessment of a community should be based on accepted ecological concepts and allow quantification of wildlife community features.

Aerial photography of the study area was available from 1956, before major construction of levees and significant modification of the floodplain. Workshop participants assumed that richness and abundance of wildlife species in 1956 are an acceptable standard of optimum conditions. Workshop participants agreed that the model output should be a value from 0 to 1.0, with 1.0 representing species richness and abundance patterns of riparian-dependent wildlife in 1956. It was further assumed that measurements of the vegetation and physical features of the riparian habitat could be compared with conditions in 1956 and used to predict changes in the wildlife community.

The model is driven mostly by variables whose values can be obtained from maps and aerial photographs. The model does not specifically address the habitat needs of species of special regional or national concern, such as the Snake River cutthroat trout (*Oncorhynchus clarki pleuriticus*), trumpeter swan (*Cygnus buccinator*), and bald eagle (*Haliaeetus leucocephalus*). Assessment of habitat conditions for species of special concern should be addressed separately from a general evaluation of the riparian wildlife community.

Three major criteria are assumed to influence species richness of wildlife in the study area: (1) changes in complexity and area of plant associations, particularly conversion of wetland cover types to more xeric plant communities from alterations in hydrologic patterns in the floodplain; (2) changes in physical complexity of the river channel and floodplain from alterations in the hydrologic regime; and (3) access to and connectivity of plant associations that affect dispersal or migration of wildlife in the floodplain. Each of these criteria is described in detail in the following pages. The first major section presents information from the literature on use of western riparian habitats by wildlife. The goal of the workshop was to use existing information for the model and to develop the model in a short time. Therefore, much of the information is from outside the study area, but the general concepts are assumed to be applicable to the Snake River. The second major section presents the habitat suitability index (HSI) model for estimating the

quality of the riparian habitat and the logic behind its development.

The workshop was conducted by the authors. Workshop participants included steering committee members of the Jackson Hole Flood Protection Project and biologists familiar with resource issues in the area, namely, F. Goodsell, Bureau of Reclamation; M. Whitfield, Greater Yellowstone Coalition; L. Glenn, W. Rigsby, and S. van Gykabeek, Trout Unlimited; T. Collins, J. Kiefling, D. Moody, and B. Oakleaf, Wyoming Game and Fish Department; L. Mettler, U.S. Army Corps of Engineers; A. Anderson, V. Moran, and S. Oddan, Service; and J. Kremer, U.S. Soil Conservation Service.

Characteristics of Riparian Wildlife Habitat

Riparian systems are the interface between aquatic and upland ecosystems and are characterized by distinct vegetation and fauna, high productivity, and high densities and diversity of wildlife species (Mitsch and Gosselink 1986). Riparian ecosystems in the intermountain west have a wide variety of plant communities because of elevational, geomorphic, and climatic influences (Brinson et al. 1981; Johnson and Lowe 1985). Attributes that augment the diversity of wildlife species and productivity of riparian communities include predominance of woody plant species, high soil moisture and surface water, diversity and interspersed physical features of habitat and plant communities, and corridors that facilitate animal dispersal and migration (Brinson et al. 1981).

Because of their unique attributes and relative rarity and the declining diversity of their fauna and flora, riparian communities have been recognized on a regional and national level (Johnson and Carothers 1982; Szaro and Rinne 1988; Flather and Hoekstra 1989). More than 70% of the riparian areas in the United States have been altered (Brinson et al. 1981), and the average annual loss of forested wetlands in the Rocky Mountain states is about 1% (Cooper and Lee 1987). This annual rate of loss is significant because of the scarcity of western riparian ecosystems and the importance of their ecological functions and their effect on local and regional species diversity of wildlife (Warren

and Schwalbe 1985). The Fish and Wildlife Coordination Act report for the Snake River Levee Project (U.S. Fish and Wildlife Service 1990) notes the occurrence of 192 vertebrate species in the cottonwood (*Populus* spp.)—riparian and associated wetland habitat types in the vicinity of Jackson, Wyoming.

Cumulative effects of human activities in and adjacent to wetlands have seriously reduced the quality and quantity of wetlands (Brown 1989). Elimination and degradation of riparian ecosystems and corresponding losses of wildlife habitat can be attributed to channel alteration, ground water pumping, surface water diversion, impoundment, direct removal of riparian vegetation, alteration of flooding regimes, and urbanization (Brinson et al. 1981). Contaminants, recreation, grazing, and habitat fragmentation also degrade viable habitat for wildlife in riparian systems.

Factors of Wildlife Diversity in Riparian Habitats

Factors of habitat diversity and the ability of riparian communities to support ecologically diverse and productive wildlife communities include horizontal and vertical diversity of size, shape, and spatial distribution of vegetation types and successional stages; physical complexity of the channel and floodplain; and spatial relations in the riparian plant community and the drainage system. Diminished habitat quality in riparian communities stems from at least two major causes: loss of diversity in biotic or physical composition and isolation or fragmentation.

Structure and Abundance of Riparian Vegetation

Diversity and productivity of plant communities adjacent to riverine channels are largely influenced by natural fluctuations in flow regime, which influence the transport of nutrients and successional composition of vegetation in floodplains (Ohmart et al. 1977; Fredrickson 1979; Brinson et al. 1981). Richness and abundance of wildlife seem to be more dependent on the structure and diversity

of plant communities than on the presence of particular associations of plant species (Szaro 1980; Brinson et al. 1981; Raedeke et al. 1988). The cover types in the model (Table 1) are based primarily on differences in vegetation structure and were developed by the tri-agency team. Foliage density, height, and patchiness of vegetation affect habitat complexity and species diversity and richness of wildlife in a riparian community (Brinson et al. 1981). Reductions in plant cover typically eliminate availability and quality of concealment, forage, and the density of prey species for carnivores (Raedeke et al. 1988). Reduced complexity in foliage height or replacement of multi-aged with even-aged stands of vegetation reduces complexity of vertical structure per unit area, resulting in concurrent loss of habitat diversity and diminished wildlife species richness (Brinson et al. 1981).

Horizontal diversity between stands influences interstand complexity and wildlife species richness in riparian communities. Distinct edges between communities and greater numbers of plant strata are associated with greater maturity of the vegetation complex in the riparian zone and greater diversity of habitats (Minshall et al. 1989). Diverse associations of species and age classes provide additional niches and strata per unit area for wildlife. In contrast, large uniform stands of the same age or monotypic species composition support lower richness of wildlife species per unit area.

Use of riparian habitat by birds has received more attention than use of it by other species groups (Brinson et al. 1981). No other habitat associations in North America are believed to be as important as riparian wetlands to noncolonial nesting birds (Carothers and Johnson 1975). Riparian ecosystems are critically important to non-breeding and migrant bird populations as well. Nationwide, more than 250 species of birds use riparian habitats (Szaro 1980; Brinson et al. 1981). Of the 192 vertebrate species in the riparian habitat off the Snake River, 150 (78%) are birds (U.S. Fish and Wildlife Service 1990). Of these 150 species, approximately 75% are passerines (songbirds).

Factors of use of riparian habitats by passerine birds include the quality of adjacent habitats, structural diversity, species composition of vegetation, and location of habitat (Stevens et al. 1977). Plant structure reflects resource distribution, which affects abundance, diversity, and distribu-

Table 1. Vegetation classification scheme for the Snake River near Jackson, Wyoming. Terminology generally follows Cowardin et al. (1979).

System	Assigned code	Subsystem	Class	Subclass and dominance type
Riverine	R2AB3	Lower perennial	Aquatic bed	Rooted vascular
	R3AB3	Upper perennial	Aquatic bed	Rooted vascular
	R3US1	Upper perennial	Unconsolidated shore	Cobble-gravel
	R3UB1	Upper perennial	Unconsolidated bottom	Cobble-gravel
Palustrine	PAB3	Palustrine	Aquatic bed	Rooted vascular
	PUB	Palustrine	Unconsolidated bottom	
	PEM1 ^a	Palustrine	Emergent	Persistent
	PSS1	Palustrine	Scrub-shrub	Broad-leaved deciduous
	PF01	Palustrine	Forested	Broad-leaved deciduous
Riparian	MG ^b	Riparian	Grassland	
	MS	Riparian	Shrubland	
	MFA	Riparian	Forested	Aspen >30%, <30% other
	MFALP ^c	Riparian	Forested	Aspen or lodgepole pine >30%, <30% other
	MFC ^c	Riparian	Forested	Cottonwood >30%, <30% other
	MFCA ^c	Riparian	Forested	Cottonwood or aspen >30%, <30% other
	MFCS ^c	Riparian	Forested	Cottonwood or spruce >30%, <30% other
	MFLP ^c	Riparian	Forested	Lodgepole pine >30%, <30% other
	MFS ^c	Riparian	Forested	Spruce >30%, <30% other
	MFW ^c	Riparian	Forested	Willow >30%, <30% other
Upland	N ^b			

^a Water regime modifiers for palustrine emergent vegetation classes: C = seasonally flooded (wet meadows), F = semipermanently flooded (cattail, bulrush, etc.), M = permanently flooded (open water).

^b Modifiers to upland and riparian grassland: r = residential, s = sagebrush. Modifiers to all classes: b = beavers, f = farmed, h = diked, x = excavated.

^c These vegetation classes must be followed by modifiers for height and coverage: height class 1 = 6.1–12.2 m (20–40 feet), class 2 >12.2 m (>40 feet); coverage class 1 = 30–70%, class 2 >70%.

tion of birds. Stauffer and Best (1980) concluded that vertical stratification of vegetation, sapling or tree size, plant species richness, and the presence of special habitat features (e.g., snags) were the habitat features most frequently related to abundance of bird species in riparian habitat.

Raptors and other predatory species meet many of their needs in riparian communities, which have comparatively high populations of prey and unique habitat features (e.g., roost and nest sites, protective cover; Lee et al. 1987; Lingle 1989). Several raptors, including ospreys (*Pandion haliaeetus*) and bald eagles, depend on riparian habitats for reproduction, wintering, and stopovers during migration (Vahle et al. 1986; Hunter et al. 1987).

Landscape and riparian features important to raptors should be considered in conjunction with site-specific characteristics including the quality of surrounding habitat types and the importance of riparian zones as corridors between vegetation types (Lee et al. 1987). Structurally complex systems provide maximum net productivity crucial to raptors and their prey base in the western states.

Community structure and abundance of small mammals seem to be strongly related to habitat diversity (Geier and Best 1980; Moulton et al. 1981; Racey and Euler 1982). Whereas microhabitat (e.g., physical characteristics in an individual home range) structure and heterogeneity affect habitat selection and density of small mammals,

macrohabitat (e.g., riparian communities) affects stability and community composition of small mammal populations (Adler 1988). Diversity of small mammal populations is increased through maintenance or enhancement of structurally diverse macrohabitats.

The specific needs of medium to large mammalian species in relation to riparian communities have received comparatively little attention, and knowledge of the value of these communities is largely unquantified (Ohmart and Anderson 1986; Raedeke et al. 1988). Characteristics of riparian systems important to large mammals include abundance and concentration of prey species, productivity and diversity of vegetation, early spring availability of forage, reduced snow accumulations, surface water and associated aquatic habitats, and the linear continuity of these habitats that facilitates movement and migration by animals (Jonkel and Cowan 1971; Pederson et al. 1979; Kellyhouse 1980; Reed 1981; Harris 1984; Raedeke et al. 1988). Minks (*Mustela vison*) attain their greatest densities in riparian habitats that provide numerous potential den sites and abundant foraging cover (Melquist et al. 1981; Birks and Linn 1982). High order streams in lowlands have the greatest habitat value for large mammalian species in the Northwest, but these areas are the most threatened by development (Raedeke et al. 1988).

The value of a riparian ecosystem to large mammals partially depends on its degree of difference from adjacent upland plant communities and increases with increasing differences (Brinson et al. 1981; Raedeke et al. 1988). The most important feature of riparian habitats for larger mammals is production of food, including terrestrial and aquatic vegetation, prey, and carrion (Krausman et al. 1985; Raedeke et al. 1988). Because of high soil moisture, presence of surface water, greater productivity of alluvial soils, and high structural diversity, riparian communities generally produce more food per unit area than uplands.

Losses of riparian wildlife habitat from inundation, urbanization, and conversion to other land uses are obvious. A less conspicuous damage of wildlife habitat quality results from livestock grazing in riparian communities throughout the intermountain west (Cope 1979; Cooper and Lee 1987). No grazing system has been beneficial to wildlife

in riparian communities (Behnke 1979). Excessive concentrations of livestock in riparian ecosystems can reduce water quality, degrade channel morphology, inhibit regeneration of woody vegetation, reduce density and quality of herbaceous vegetation, and alter plant composition and structural characteristics (Cope 1979; Hogan 1979; Munther 1982; Kauffman and Krueger 1984; Kochert et al. 1986; Szaro and Rinne 1988). Lotic habitats in livestock exclosures in Colorado had narrower stream widths, greater stream depths, and riparian vegetation of greater density and diversity than grazed stream reaches (Stuber 1985). Although a grazed riparian study site in Nevada did not seem to have received excessive use by livestock, species diversity, species richness, and biomass of small mammals were all higher on a comparable ungrazed site (Medin and Clary 1989). The most obvious differences in vegetation between sites was a reduction of forb and graminoid heights and graminoid biomass by approximately 50% in the grazed site. Larger home range size and lower densities of minks were recorded in Montana where riparian communities supported lower plant density because of livestock grazing (Mitchell 1961). Numbers and species richness of birds were greater in ungrazed exclosures than in grazed stream reaches in Arizona and New Mexico (Szaro and Rinne 1988). Differences in physical and vegetative attributes between grazed and ungrazed riparian sites included greater bank stability and higher abundance of woody vegetation on ungrazed sites.

Channel and Floodplain Complexity

Channel Configuration

Multiple channels from lateral migration of the main channel and sinuosity in lotic systems contribute to habitat diversity of riverine and riparian communities. In riverine systems where large amounts of debris and sediment are dropped rapidly, channels are braided and have a high width-to-depth ratio (Heede 1980; Platts et al. 1983). Physical characteristics of braided river channels include numerous active side channels; relatively flat, shallow beds; heavy sediment load; erodible banks; and frequent and rapid variation in discharge.

The consequences of stream braiding include increased surface area of banks, greater potential for accumulation of sediments and organic materials, and greater deposition of organic material in the aquatic system from adjacent riparian vegetation (McArthur 1989). Biological diversity is greater in habitats of a braided stream than in habitats of a nonbraided stream of comparable size because the horizontal dimension of a braided stream is larger and therefore promotes a more diverse food base (Brinson et al. 1981; McArthur 1989). Erosion and deposition of sediments associated with braided or meandering channels have physical features such as meander bends, point bars, undercut banks, vegetated banks, and diversity in bed material that provide microhabitats for feeding and breeding by numerous aquatic and wetland-dependent wildlife species. In contrast, constructed channels typically reduce stability of stream banks, lower diversity of microhabitats, reduce abundance or completely eliminate vegetation at the interface of land and water, and increase loads of suspended solids.

Diversity of wildlife species in riparian communities is enhanced by the presence of wetland and open-water cover types in addition to the lotic environment (Fredrickson 1978; Brinson et al. 1981; Sedell and Froggatt 1984; Holden et al. 1986). Wetland cover types (e.g., oxbows, seasonally flooded wetlands, spring creeks) in close association with the riverine channel enhance structural and vegetation diversity.

Dynamically Stable Islands

As a consequence of lateral movement of the main channel, side channels are developed that create and maintain islands throughout the floodplain. Diversity in the structure and composition of plant communities on in-channel islands provides diversity in the structure and vegetation of habitats in the floodplain and in the channel. Because islands and their associated plant communities are alternately destroyed and created, they provide a relatively stable area and diversity of habitats. Seasonally high discharges in channelized reaches or levees increase erosion of these islands and decrease habitat diversity in modified reaches of the floodplain. Although sediment deposition in modified reaches may provide suitable substrates for establishment of early seral stages

of wetland vegetation, seasonally high discharges prevent long-term persistence of islands.

Overbank Flooding

Natural levees (from seasonal increases in discharge), overbank flooding, and sediment deposition generally support a high diversity of various wetland types and vegetation communities in riverine floodplains. Natural levees tend to regulate the rate at which river water is distributed across the floodplain during flooding (Minshall et al. 1989). Modification of water regimes throughout the floodplain may significantly influence the structure of riparian communities through seasonal reduction in surface water and deposition of organic debris and through alteration in complexity and diversity of wetland habitats. Stabilized or static water regimes, which are either wetter or drier than natural conditions (Fredrickson 1980), eliminate or greatly modify plant and animal associations tied to fluctuating water regimes. Longer flood duration and greater water depth are characteristic of riverine channels in levees. Conversely, elimination of seasonal flooding of riparian communities results in vegetation and faunal shifts away from wetland-associated species (Martin 1977; Ohmart et al. 1977; Mitsch and Gosselink 1986). Reduction or elimination of peak flows throughout the floodplain can eventually lower the water table, which causes replacement of wetland-associated communities with vegetation of more xeric conditions.

Shoreline Complexity

Partial or total removal of streamside vegetation diminishes diversity and quality of riparian habitat and increases sediment loads, which reduces bed-form roughness and thereby decreases lotic habitat diversity (Platts et al. 1985). The importance of debris in riverine channels is in its effect on channel hydraulics and ultimately on populations of prey species (Angermeier and Karr 1984; Sedell and Swanson 1984; Bisson et al. 1987). The orientation and size of woody debris can enhance channel and habitat heterogeneity by altering flow direction and velocity (Everest and Meehan 1981). Deposited organic matter provides structure and stability for food resources in most riverine systems (McArthur 1989). Therefore, alterations that limit or remove organic matter or significantly change the nature of these materials affect the productivity, diversity,

and stability of aquatic communities. Lower diversity and abundance of strata (e.g., rocks, large woody debris) in and adjacent to flowing water reduce (1) the accumulation and decomposition of organic debris, resulting in decreased production of macroinvertebrates, (2) production of invertebrate and fish biomass by elimination of cover and disruption of orientation and territorial behavior, and (3) prey availability for piscivorous species (Marzolf 1978). Habitat quality for wetland-associated furbearers is degraded when snags, boulders, aquatic and riparian vegetation, and shoreline complexity are reduced (Arner et al. 1976; Gray and Arner 1977; Racey and Euler 1983).

Investigations in riparian communities indicate that abundance and diversity of riparian herpetofauna are dependent on the availability and diversity of microhabitats in large woody debris and litter provided, in part, by seasonal flooding (Jones and Glinski 1985; Jones 1988). Although rapidly fluctuating water levels may be temporarily detrimental to riparian herpetofauna (Warren and Schwalbe 1985), debris deposited by high flows provides microhabitats beneficial to species not immediately dependent on the aquatic portion of the riparian community. Because of reductions in seasonal flooding, density of woody vegetation, and abundance of debris, species richness and abundance of herpetofauna are lower in regulated stream reaches than in reaches with natural fluctuations in discharge.

Debris contributes to habitat structure in the channel by anchoring the position of pools and creating backwaters along the stream boundary and by contributing to lateral migration of the channel (Sedell and Fraggate 1984; Bisson et al. 1987). In higher order streams or in rivers with channels that are too wide for spanning, debris is deposited along channel margins. The highest densities of invertebrates and the most productive fish habitat are in these locations. These sites provide excellent foraging habitat for minks and other wetland-associated furbearers because debris provides cover for prey as well as for hunting predators. Woody debris has a similar influence on the quality of habitat and its use by river otters (*Lutra canadensis*; Melquist and Hornocker 1983). Availability of prey was believed to have the greatest influence on habitat use by river otters in Idaho. However, adequate shelter was required for exten-

sive use of foraging areas. Because they provided ideal cover for foraging and resting, logjams and debris were frequent resting sites and centers of activity for river otters.

Spring-fed Creeks

Spring-fed tributary creeks are an important wetland class in the Snake River floodplain. They provide suitable temperatures and essential spawning habitat for cutthroat trout and other fish species (Kiefling 1978; Simpson and Wallace 1982). The importance and rarity of suitable spawning habitat for cutthroat trout in the upper Snake River drainage was emphasized by Kiefling (1978). Spawning habitat in the main river channel is extremely limited because high flows during spring runoff increase turbidity and sediment loads during the spawning period.

Habitat in spring-fed tributaries on the Snake River floodplain is believed to have deteriorated in area and quality from alterations in seasonal flooding and sedimentation (U.S. Fish and Wildlife Service 1990). These factors in combination with decreased discharge of some spring-fed creeks reduced habitat for all life stages of trout. Conversely, sedimentation and plant succession in spring-fed creeks have been beneficial to other species (e.g., trumpeter swan) by providing wetland cover types suitable for foraging and reproduction. Total dewatering or absence of spring-fed creeks in the floodplain of a given reach of the Snake River is assumed to create lower habitat diversity and a potentially lower richness of wildlife species.

Pool-Riffle Ratio

Pool-riffle ratio is the length or percentage of riffle divided into the length or percentage of pool (Platts et al. 1983). Pools are the deeper, placid, and slower moving sections of a river or stream, whereas riffles are the faster, shallower waters. A ratio of 1:1 is generally believed to provide optimum conditions for fish production and habitat quality (Platts et al. 1983; Marcus et al. 1990). A well-interspersed mix of pools and riffles provides a variety of microhabitats from which numerous aquatic and wetland dependent species benefit. Physical obstruction to water flow increases complexity and productivity of lotic communities by contributing to formation of pools, storage of sediment and organic matter, and provision of cover for

invertebrates and vertebrates. Obstructions create pools with deeper water of slower velocity that provide escape cover and promote higher densities of fishes in various age classes (Angermeier and Karr 1984; Sedell and Froggatt 1984; Bisson et al. 1987).

Spatial Relations

The ecological value of wetlands is largely based on their interspersed and physical relation with other plant communities (Mitsch and Gosselink 1986). Resident wildlife species with specific, unique habitat requirements usually have relatively small home ranges that may be within a single wetland community. In contrast, large, wide-ranging species are not restricted to riparian communities, but may be only seasonally dependent on them. Nevertheless, access to and use of riparian communities may be essential to meeting annual habitat requirements of these species.

Landscape modifications that eliminate movement between component subsystems may be as devastating as direct elimination of required critical habitats (Harris 1988). For example, riparian corridors are believed to be important for dispersal of small mammals (Cross 1985; Dickson and Williamson 1988). Ideally, riparian communities are evaluated and managed for their role in a functional landscape mosaic, large enough to provide significant wildlife habitat and capable of resisting detriment from adjacent lands (Harris 1988; Olson and Knopf 1988; Brown 1989). Urbanization and intensive land use tend to produce isolated units of wetland habitat where wildlife and wetland functions face higher susceptibility and lower resiliency to detrimental effects from surrounding land use (Brown 1989). Although riparian communities can be considered biological islands that support fauna and flora of greater species richness and densities than are typical of adjacent communities, the integrity of wildlife values in riparian wetlands cannot be maintained by setting aside reserves and disregarding landscape-level processes.

Corridor width and breaks in the corridor are critical influences of habitat value in a stream corridor (Forman 1983). Fragmentation and elimination of access to critical habitats result in loss of wide-ranging species, reduction of population vi-

ability of area-sensitive and interior species, reduction in genetic integrity of species or populations, and enhancement of habitat quality for generalist species of disturbed environments (Brinson et al. 1981; Cooper and Lee 1987). The final result is that riparian communities, and eventually regions, lose distinguishing biological characteristics and acquire already common generalist species (Sampson and Knopf 1982; DeGraff 1986; Harris 1988).

Most forms of landscape degradation can be attributed to reduction in landscape complexity that lowers diversity of plant and animal communities (Bridgewater 1988). The role of riparian habitats in maintenance of regional wildlife values must be addressed if integrity of regional ecosystems is to be maintained (Knopf 1985; Olson and Knopf 1988). Within-habitat (alpha) diversity has received widespread use in wildlife management and in most situations can be expected to be high in structurally diverse riparian communities. However, most measures of alpha diversity do not address the spatial distribution of species or the composition of entire communities. The potential for high levels of wildlife species richness is greater between than within plant communities. Because of their unique attributes, riparian communities of western riverine systems contribute to between habitat (beta) diversity and regional (gamma) diversity. Greater emphasis needs to be placed on techniques that address size, shape, distribution, and linkages of vegetative communities in large geographic areas if wildlife values in riparian ecosystems are to be preserved or enhanced (Odum 1978; Sanderson et al. 1979; Gall and Christian 1984; Harris 1988).

Summary of Riparian Habitat and Wildlife Diversity

A common theme in describing the quality of riparian plant communities as wildlife habitat is physical, vegetative, and spatial diversity. The ability of riparian areas to support diverse and productive wildlife communities is a function of the physical complexity of the water channel and immediate shoreline and of the abundance and diversity of plant communities adjacent to the riverine channel. Actions that modify naturally fluctuating

water regimes, eliminate or reduce channel and shoreline diversity, diminish vertical or horizontal vegetative structural complexity, or fragment and isolate riparian communities tend to lower habitat quality for species that are restricted to or are seasonally dependent on riparian habitats.

The role of western riparian wetlands in regional wildlife diversity is less well defined than characteristics of these communities to specific wildlife species. On a landscape level, the role of riparian corridors in dispersal and migration of mammals and birds alone justifies greater protection and enhancement of the unique attributes of these communities. The value of riparian communities to wildlife increases in response to greater divergence in vegetative characteristics between riparian and adjacent vegetative communities. As a result, riparian communities along high order, low elevation riverine cover types probably have the greatest values for wildlife in the intermountain and southwestern United States.

Habitat Suitability Index Model for a Riparian Wildlife Community

Model Overview and Output

This model is designed for a quantitative assessment of the quality of the riparian habitat along the Snake River near Jackson, Wyoming. The model estimates the richness of vertebrate species that are dependent on riparian habitat by comparing vegetation and physical conditions with those that were present along the river in 1956. The assumption is that the wildlife community in 1956 was highly diverse and its abundance patterns approximate historical norms. The output of the model is a measure of the richness of vertebrates that depend on riparian habitat and is based on the assumption that as it increases, richness increases and the pattern of species abundance more closely approximates the 1956 baseline conditions. The quality of riparian habitat can be evaluated for discrete 1.6-km river reaches or for the entire 40 km area. The list of riparian-dependent species (Appendix) was determined from a rating system by Gerhart and Olson (1982).

Model Applicability

Application of the model is restricted to the Snake River near Jackson, Wyoming. The model relies in part on a specific comparison of existing or predicted plant communities with conditions in 1956.

Model Variables and Assumptions

The following sections describe the specific variables for estimating vertebrate species richness with the model. These sections explain the derivation of the variables, including interpretation of the scientific literature and incorporation of workshop results. The model is intended to be applied with remotely sensed data describing the model variables. Adequate data for an empirically based statistical model do not exist.

SIVI—Percentage of 1956 Cover Type

The scientific literature and the workshop participants support the hypothesis that the horizontal and vertical structure provided by a diversity of plant community types and successional stages is the primary factor influencing wildlife species richness in riparian habitats. Workshop participants were concerned that a model output equal to a general measure of species richness might overlook the habitat needs of species with specific needs (e.g., species requiring large stands of mature cottonwood). Therefore, the model is based on comparing existing or expected vegetation patterns with conditions in 1956, and yields existing or expected richness of riparian-dependent species from this comparison. Such a measure avoids the pitfalls of managing for maximum alpha (site-specific) species richness, but should address the needs of species important in maintaining gamma (regional) species richness.

The model variable that assesses vegetation structure is the existing or expected vegetation by cover type expressed as a percentage of the vegetation in 1956. The cover types that are mapped and used for this variable to assess habitat on the Snake River near Jackson, Wyoming, are listed in Table 1. This cover type system was developed by the U.S. Army Corps of Engineers in cooperation with other members of the tri-agency team. The

successional stage of woody vegetation is considered in the modifiers for cottonwood stands and the cover types for cottonwood or willow shrub. Future years with lower than 1956 amounts of any particular successional cover type will cause the model to yield lower values of species richness. Each cover type in 1956 is assumed to have made an important contribution to species richness in 1956.

The curve of the suitability index (SI) graph for this variable (SIV1; Fig. 1) is based on species-area relation concepts in community ecology. A general guiding principle in community ecology is that, as the area of a habitat is reduced to one-tenth its original size, the number of species is reduced to one-half (Wilson 1985). The specific shape of this curve is based on species-area relations for birds from 34 shelterbelts in Kansas (Schroeder et al. 1992). Empirical data on species-area relations in Wyoming riparian habitat patches are limited. Gutzwiller and Anderson (1987) provided data on riparian habitat fragments and use by 10 species of cavity nesting birds. A species-area curve with their data did not differ from a species-area curve for cavity nesters in the Kansas shelterbelt data ($P < 0.05$). Therefore, the shape of the species-area curve from the Kansas data is assumed to be adequate for use in this model.




Fig. 1. Suitability index (SIV1)—the relation between the percentage of 1956 cover type and the SI value, determined as: $SI = 0.194 \times (\% \text{ of } 1956)^{0.357}$. If the observed cover exceeds 100%, the SI should be set at 1.0.

The shape of the graph in SIV1 indicates that the rate of change in SI values is low for cover types with only small deviations from 1956 cover type percentages and much higher in cover types with extremely large deviations from 1956. Thus, a 20% deviation from 1956, from 100% similarity to 80%, results in an estimated 7% decline in species richness. A similar 20% decline, from 30% to 10% similarity, however, results in an estimated 21% decline in richness. This relation indicates that, although 1956 is the desired condition, minor deviations cause only small changes in the model output. This reflects a concern that percentages of various cover types in a naturally functioning dynamic riparian system vary over time and that only as these differ grossly from 1956 is there a severe effect on species richness. Cover types exceeding the areas in 1956 cannot exceed a 1.0 SI. The total area of each river reach in 1956 (to the outer boundaries of the estimated 500-year floodplain) is used in all computations for other years, irrespective of changes in cover type composition or conversion to upland or urban habitats.

An important aspect of comparing vegetation patterns with 1956 baseline data is the spatial distribution of vegetation along the entire 40-km study area. For purposes of assessing this linear distribution, the variable SIV1 should be applied in each 1.6-km river segment for all cover types. The 1.6-km distance was chosen because the study area had previously been divided into these segments. Workshop participants agreed that, although the distance was somewhat arbitrary, it seemed to be biologically reasonable, given the small home range size of most of the riparian-dependent species of wildlife.

The procedure for computing a suitability index value for this variable (SIV1) involves the following steps:

1. Determine the relative percent abundance of each plant cover type in each river segment for the 1956 baseline condition:

$$\frac{\text{Area of cover type}}{\text{Area of river segment}} = \% \text{ of cover type (1956)}$$

2. For the desired year of analysis, determine the relative percent abundance of each cover type in each river segment:

$$\frac{\text{Area of cover type}}{\text{Area of river segment}} = \% \text{ of cover type (year of analysis)}$$

3. Compare the percent occurrence of each cover type for the desired year of analysis with the percent 1956 baseline for that cover type. Convert this to a percentage:

$$\frac{\% \text{ cover type (year of analysis)}}{\% \text{ cover type (1956)}} \times 100 = \frac{\% \text{ of 1956 (for year of analysis)}}{\% \text{ of 1956 (for year of analysis)}}$$

4. Put the percent of 1956 (for year of analysis) for each cover type into the formula for the graph for SIV1 and determine the SI value for each cover type.
5. Determine an area-weighted SI by multiplying the value for SIV1 for each cover type by the decimal equivalent of the percentage cover type (1956; determined in step 1) for that cover type. (For example, if a cover type had 50% abundance in 1956, use the decimal 0.5.)
6. Sum the values from step 5 for all cover types in the river segment being analyzed. This sum is the overall value for SIV1 for this segment.

For further details on applying this variable, refer to the sample data sets and calculations in the section *Habitat Suitability Index Determination and Application of the Model*.

SIV2—Channel and Floodplain Complexity

The workshop participants and pertinent literature support inclusion of a variable to assess physical complexity of the channel and floodplain for assessment of habitat quality in riparian communities. Factors deemed important by workshop participants included the presence or absence of overbank flooding, sediment-free spring-fed creeks, physically complex shoreline, high pool-riffle ratios, braided channels, oxbows, and islands. All of these factors contribute to defining a river system that is dynamic and influenced by fluctuations in discharge and seasonal distribution of water throughout the floodplain.

The suitability index for this variable (SIV2) is determined by summing a single value for each of the six factors of channel and floodplain diversity.

Factor	Index
1. Channel configuration	
a. main channel braided or meanders throughout erosional floodplain; oxbows, side channels, or other nonriverine wetland types are present in the evaluated reach	0.2
b. single channel; reach is relatively straight; portions of the channel are contained in dikes to prevent lateral movement of main channel in floodplain; other wetland types not present in the evaluated reach	0.1
c. entire reach is channelized; river is confined on both sides to prevent lateral movement in floodplain	0.0
2. Dynamically stable islands	
a. islands are present within annual boundaries of main channel; islands of various size supporting full spectrum of vegetation succession ranging from newly deposited, unvegetated sediment to islands dominated by mature stands of riparian vegetation (e.g., willow and cottonwood)	0.2
b. islands are present within annual boundaries of main channel, vegetation communities on islands are skewed to one end of successional spectrum (e.g., majority in mature class or composed of newly deposited, unvegetated sediment)	0.1
c. islands absent in the evaluated reach	0.0
3. Overbank flooding	
a. high potential for seasonal overbank flooding, channel not confined by artificial barriers (e.g., dikes and levees)	0.2
b. moderate potential for seasonal overbank flooding; one side or segments of channel confined by dikes or levees	0.1
c. little to no potential for seasonal overbank flooding; entire channel in evaluated reach is channelized	0.0
4. Spring-fed creeks	
a. present, diverse substrate composition	0.2
b. present, more homogenous substrate composition	0.1
c. absent	0.0
5. Shoreline complexity	
a. presence of undercut banks, large woody debris (e.g., root wads, tree trunks, and debris piles) im-	

mediately adjacent to channel and in the water column; vegetated banks in immediate proximity to water; diversity of substrate types (e.g., silt, sandbars, gravel, cobble, and boulders) compose land-water interface

- b. shoreline essentially straight, open and exposed (i.e., channelized and levee) with abrupt monotypic edge between water and land; little to no diversity in substrates at land-water interface; few to no physical or vegetation features that provide structure or cover in or in close to water 0.1
6. Pool-riffle ratio 0.0
- a. good, area in pools and riffles approximately equal 0.1
- b. poor, highly uneven distribution of either pools or riffles-runs 0.0
- Total score (0-1.0) ____

These factors and their perceived importance to the wildlife community were weighted by participants of the workshop. Although some degree of intercorrelation of the above factors is probable, the participants decided to include all factors to assure an adequate overall view of physical complexity. This variable should be applied by computing an SI value for each river segment and averaging them for the entire study area. The average SIV2 for 1956 conditions is assumed to be the best attainable condition, and overall average conditions following 1956 and the influence of the levees should always be of lower quality. Suitability index values for an individual river segment could exceed the SI of that segment for 1956. The overall average condition of channel and floodplain complexity is assumed to be more important than a spatial distribution of these features equal to 1956.

SIV3—Percentage of River Reach Subject to Human Disturbance

The ability of riparian wildlife species to move freely and disperse along the riparian corridor is important to the long-term maintenance of wildlife species richness. Workshop participants and the scientific literature indicate that excessive

human disturbance can have a negative influence on wildlife in riparian habitats.

Human disturbance is assumed to have two major effects: (1) direct loss of use of habitat in and near areas of disturbance, and (2) disruption of movement patterns of wildlife along the narrow riparian corridor. The effects of human disturbance on wildlife are difficult to quantify. Certain species and, in some cases, certain individuals in a species, respond differently to human disturbances. The effects of human disturbances often extend beyond the boundaries of the area where the disturbance is effected. We assume that disturbances are in one of three categories and that more severe disturbances affect habitat use and wildlife movement in a larger surrounding zone as delineated in Table 2.

The distances provided for the additional zone of influence are based on data about disturbed bald eagles (McGarigal et al. 1991). A buffer zone of approximately 100 m (328 feet) was estimated to protect 50% of the breeding eagle population from flushing by human disturbances. The suitability index for assessing the effects of human disturbance (SIV3) is illustrated in Fig. 2. The percent of the disturbed river reach is determined by combining the actual disturbed area and the area in the surrounding zone of influence for each disturbance and computing a percentage of the disturbed reach. As this percentage increases, the SI value is assumed to linearly decrease, reaching

Table 2. *Level and extent of influence of human activities on wildlife habitat.*

Level of disturbance	Examples of activities or land uses	Additional zone of influence
High	Industry, commercial development, high density residential (>1 unit per 1.2 ha) development, paved highway, active quarry	100 m
Medium	Levee with public access, golf course, low density residential (<1 unit per 1.2 ha) development, gravel road, inactive quarry, cropland	50 m
Low	Dirt road, levee with limited access, recreation areas, rangeland, native pasture	0 m




Fig. 2. Suitability index (SIV3)—the relation between the percentage of disturbed river reach and the SI value.

a worst case value of 0.2 at 100%. Twenty percent of the riparian dependent native species (Appendix) are assumed to tolerate a high level of human disturbance and continue to occupy highly disturbed river reaches.

Habitat Suitability Index Determination and Application of the Model

Application of the model requires the computation of an intermediate value for SIV1, SIV2, and SIV3 along each 1.6-km river reach segment. Overall HSI determination then uses the averages of the SI values for all reaches. Separate HSI values are not computed for each river reach. Vegetation structure is assumed to be a more important determinant of species richness than channel complexity. Therefore, SIV1 is weighted in the HSI formula. The overall HSI for the entire study area is computed as follows:

$$\text{Study area HSI} = \frac{2(\text{SIV1}) + (\text{SIV2})}{3} \times (\text{SIV3})$$

The following sample data set illustrates how to apply the model and calculate river segment SIs

and the overall HSI. Any modifications of the original model should be accompanied by further analysis of sample data sets to ensure appropriate response in the model.

SIV1—Structure and Abundance of Riparian Vegetation.

Step 1. Determine relative percentage of cover types for 1956.

Cover type	Assumed 1956 area (ha)	Relative %
Riparian cottonwood (>40 feet, >70%)	100	50
Palustrine scrub-shrub wetland, willow	60	30
Palustrine emergent wetland	20	10
Riverine	20	10
Total	200	100

Step 2. Determine relative percentage of cover types for year of analysis.

Cover type	Year of analysis area (ha)	Relative %
Riparian cottonwood (>40 feet, >70%)	40	20
Palustrine scrub-shrub wetland, willow	40	20
Palustrine emergent wetland	10	5
Riverine	20	10
Upland grassland	90	45
Total	200	100

Step 3. Compare year of analysis with 1956.

Cover type	Year of analysis %	1956 %	% of 1956
Riparian cottonwood	20	50	40
Palustrine scrub-shrub wetland, willow	20	30	67
Palustrine emergent wetland	5	10	50
Riverine	10	10	100
Upland grassland ^a	45	0	NA

^a Upland grassland is not considered because it did not occur in the river reach in 1956.

Step 4. Determine SIV1 from suitability index graph and formula. Put percentage of 1956 cover type (use the whole number) in the formula and calculate SI.

Cover type	SIV1
Riparian cottonwood	0.72
Palustrine scrub-shrub wetland, willow	0.87
Palustrine emergent wetland	0.78
Riverine	1.00
Upland grassland	NA

Step 5. Multiply the SI for each cover type by the percentage of 1956.

Cover type	SIV1	1956 %	Weighted SI
Riparian cottonwood	0.72	× 50	= 0.36
Palustrine scrub-shrub wetland, willow	0.87	× 30	= 0.26
Palustrine emergent wetland	0.78	× 10	= 0.078
Riverine	1.00	× 10	= 0.1
Upland grassland	NA		

Step 6. Determine the overall value for SIV1 in the river segment.

Sum of weighted SIs from step 5 = SIV1 = 0.798

SIV2—Channel and Floodplain Complexity

Factor	River segment Index	score
1. Channel configuration		
a. main channel braided or meanders throughout erosional floodplain, oxbows, or side channel or other nonriverine wetland types are present in reach being evaluated	0.2	
b. single channel reach is relatively straight, portions of the channel are contained in dikes to prevent lateral movement of main channel in floodplain, other wetland types not present in reach being evaluated	0.1	
c. entire reach is channelized, river confined on both sides to		

prevent lateral movement in the floodplain

0.0 0.0

2. Dynamically stable islands

a. islands are present in annual boundaries of main channel, islands of various size supporting full spectrum of vegetation succession range from newly deposited, unvegetated sediment to islands dominated by mature stands of riparian vegetation (e.g., willow, cottonwood)

0.2

b. islands are present in annual boundaries of main channel; vegetation communities on islands are skewed to one end of successional spectrum (e.g., majority in mature class or composed of newly deposited, unvegetated sediment)

0.1

c. islands absent in the evaluated reach

0.0 0.2

3. Overbank flooding

a. high potential for seasonal overbank flooding, channel not confined by artificial barriers (e.g., dikes, levees)

0.2

b. moderate potential for seasonal overbank flooding, one side, or segments or channel confined by dikes or levees

0.1

c. little to no potential for seasonal overbank flooding; the entire channel in evaluated reach is channelized

0.0 0.0

4. Spring-fed creeks

a. present, diverse substrate composition

0.2

b. present, more homogenous substrate composition

0.1

c. absent

0.0 0.1

5. Shoreline complexity

a. presence of undercut banks, large woody debris (e.g., root

wads, tree trunks, debris piles)
immediately adjacent to
channel and in the water
column; vegetated banks in
immediate proximity to water;
diversity of substrate types
(e.g., silt, sandbars, gravel,
cobble, boulders) compose
land-water interface

0.1

- b. shoreline essentially straight,
open and exposed (i.e.,
channelized, levee) with abrupt
monotypic edge between water
and land; little to no diversity
in substrates at land-water
interface; few to no physical or
vegetation features that provide
structure or cover in or in close
proximity to water

0.0

0.0

6. Pool-riffle ratio

- a. good, area in pools and riffles
approximately equal

0.1

- b. poor, highly uneven distribution
of either pools or riffles-runs

0.0

0.0

Total score SIV2 (0-1.0) = 0.3

SIV3—Spatial Relations and Human Disturbance

	Percent SIV3	
Percentage of river reach subject to human disturbance	20	0.84

Study Area HSI

Use average values for SIV1, SIV2, and SIV3 for
all river reaches of concern. In this example, only
one reach is used.

$$\frac{2(\text{SIV1}) + (\text{SIV2})}{3} \times (\text{SIV3}) = \frac{2(0.798) + 0.3}{3} \times 0.84 = 0.53$$

The final HSI value of 0.53 indicates a riparian
habitat that supports approximately one-half the
number of riparian-dependent species as the same
area in 1956.

References

- Adler, G. H. 1988. The role of habitat structure in organizing small mammal populations and communities. Pages 289-299 in C. Szaro, K. E. Severson, and D. R. Patton, technical coordinators. Management of amphibians, reptiles, and small mammals in North America. U.S. Forest Service General Technical Report RM-166.
- Angermeier, P. L., and J. R. Karr. 1984. Relationships between woody debris and fish habitat in a small warmwater stream. Transactions of the American Fisheries Society 113:716-726.
- Arner, D. H., H. R. Robinette, J. E. Frasier, and M. H. Gray. 1976. Effects of channelization of the Luxapalila River of fish, aquatic invertebrates, water quality, and furbearers. U.S. Fish and Wildlife Service FWS/OBS-76/08. 58 pp.
- Behnke, R. J. 1979. Values and protection of riparian ecosystems. Pages 164-167 in G. A. Swanson, technical coordinator. The mitigation symposium: a national workshop on mitigating losses of fish and wildlife habitats. U.S. Forest Service General Technical Report RM-65.
- Birks, J. D. S., and I. J. Linn. 1982. Studies of home range of the feral mink (*Mustela vison*). Symposium of the Zoological Society of London 49:231-257.
- Bisson, P. A., R. E. Bilby, M. D. Bryant, D. A. Dolloff, G. B. Grette, R. A. House, M. L. Murphy, K. V. Doski, and J. R. Sedell. 1987. Large woody debris in forested streams in the Pacific Northwest: past, present and future. Pages 143-190 in E. O. Salo and T. W. Cundy, editors. Streamside management: forestry and fishery interactions. University of Washington Institute of Forest Resources Contribution 57.
- Bridgewater, P. B. 1988. Biodiversity and landscape. Earth-Science Reviews 25:485-491.
- Brinson, M. M., B. L. Swift, R. C. Plantico, and J. S. Barclay. 1981. Riparian ecosystems: their ecology and status. U.S. Fish and Wildlife Service FWS/OBS-81/17. 151 pp.
- Brown, M. T. 1989. Forested wetlands in urbanizing landscapes. Pages 19-26 in D. D. Hook and R. Lea, editors. Proceedings of the symposium: the forested wetlands of the southern United States. U.S. Forest Service General Technical Report SE-50.
- Carothers, S. W., and R. R. Johnson. 1975. Water management practices and their effects on nongame birds in range habitats. Pages 210-222 in D. R. Smith, technical coordinator. Proceedings of the symposium on management of forest and range habitats for nongame birds. U.S. Forest Service General Technical Report WO-1.

- Cooper, D. J., and L. C. Lee. 1987. Rocky Mountain wetlands: ecosystems in transition. *National Wetlands Newsletter* 9(3):2-6.
- Cope, O. B., editor. 1979. Proceedings of the forum—grazing and riparian/stream ecosystems. Trout Unlimited, Inc., Vienna, Va. 94 pp.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31. 131 pp.
- Cross, S. P. 1985. Responses of small mammals to forest riparian perturbations. Pages 269-275 in *Riparian ecosystems and their management: reconciling conflicting uses*. U.S. Forest Service General Technical Report RM-120.
- DeGraaf, R. M. 1986. Urban bird habitat relationships: application to landscape design. *Transactions of the North American Wildlife and Natural Resources Conference* 51:232-259.
- Dickson, J. G., and J. H. Williamson. 1988. Small mammals in streamside management zones in pine plantations. Pages 375-378 in R. C. Szaro, K. Severson, and D. R. Patton, technical coordinators. *Management of amphibians, reptiles, and small mammals in North America*. U.S. Forest Service General Technical Report RM-166.
- Everest, F. H., and W. R. Meehan. 1981. Forest management and anadromous fish habitat productivity. *Transactions of the North American Wildlife and Natural Resources Conference* 46:521-530.
- Flather, C. H., and T. W. Hoekstra. 1989. An analysis of the wildlife and fish situation in the United States: 1989-2040. U.S. Forest Service General Technical Report RM-178. 147 pp.
- Forman, R. T. T. 1983. Corridors in a landscape: their ecological structure and function. *Ekologia (CSSR)* 2:375-387.
- Fredrickson, L. H. 1978. Lowland hardwood wetlands: current status and habitat values for wildlife. Pages 296-306 in P. E. Greeson, J. R. Clark, and J. E. Clark, editors. *Wetland functions and values: the state of our understanding*. Proceedings of the National Symposium on Wetlands. American Water Resources Association, Minneapolis, Minn.
- Fredrickson, L. H. 1979. Floral and faunal changes in lowland hardwood forests in Missouri resulting from channelization, drainage, and impoundment. U.S. Fish and Wildlife Service FWS/OBS-78/91. 130 pp.
- Fredrickson, L. H. 1980. Management of lowland hardwood wetlands for wildlife: problems and potential. *Transactions of the North American Wildlife and Natural Resources Conference* 45:376-386.
- Gall, B. C., and K. R. Christian. 1984. A program for a wildlife inventory. *Journal of Environmental Management* 19:277-289.
- Gerhart, W. A., and R. A. Olson. 1982. Handbook for evaluating the importance of Wyoming's riparian habitat to terrestrial wildlife. Wyoming Game and Fish Department, Cheyenne. 92 pp.
- Geier, A. R., and L. B. Best. 1980. Habitat selection by small mammals of riparian communities: evaluating effects of habitat alterations. *Journal of Wildlife Management* 44:16-24.
- Gray, M. H., and D. H. Arner. 1977. The effects of channelization on furbearers and furbearer habitat. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies 31:259-265.
- Gutzwiller, K. J., and S. H. Anderson. 1987. Multi-scale associations between cavity-nesting birds and features of Wyoming streamside woodlands. *Condor* 89:534-548.
- Harris, L. D. 1984. *The fragmented forest*. University of Chicago Press, Chicago, Ill. 208 pp.
- Harris, L. D. 1988. The nature of cumulative impacts on biotic diversity of wetland vertebrates. *Environmental Management* 12:675-693.
- Heede, B. H. 1980. Stream dynamics: an overview for land managers. U.S. Forest Service General Technical Report RM-72. 26 pp.
- Hogan, G. 1979. Managing riparian ecosystems (zones) for fish and wildlife in eastern Oregon and eastern Washington. U.S. Forest Service, Portland, Oreg. 44 pp.
- Holden, P. B., R. D. Hugie, L. Crist, S. B. Chanson, and W. J. Masslich. 1986. Development of a fish and wildlife classification system for backwaters along the lower Colorado River: final report. BioWest, Inc., Logan, Utah. 207 pp.
- Hunter, W. C., R. D. Ohmart, and B. W. Anderson. 1987. Status of breeding riparian obligate birds in southwestern riverine systems. *Western Birds* 18:10-18.
- Johnson, R. R., and S. W. Carothers. 1982. Riparian habitats and recreation: interrelationships and impacts in the Southwest and Rocky Mountain region. *Eisenhower Consortium Bulletin* 12. 31 pp.
- Johnson, R. R., and C. W. Lowe. 1985. On the development of riparian ecology. Pages 112-116 in R. R. Johnson, C. D. Ziebell, D. R. Patton, P. F. Ffolliott, and R. H. Hamre, technical coordinators. *Riparian ecosystems and their management: reconciling conflicting uses*. First North American Riparian Conference: 1985. U.S. Forest Service General Technical Report RM-120.
- Jones, K. B. 1988. Comparison of herpetofaunas of a natural and altered riparian ecosystem. Pages 222-227 in R. C. Szaro, K. E. Severson, and D. R. Patton, technical coordinators. *Management of amphibians, reptiles, and small mammals in North America*. U.S. Forest Service General Technical Report RM-166.

- Jones, K. B., and P. C. Glinski. 1985. Microhabitats of lizards in a southwestern riparian community. Pages 342-346 in *Riparian ecosystems and their management: reconciling conflicting uses*. U.S. Forest Service General Technical Report RM-120.
- Jonkel, C. J., and I. M. Cowan. 1971. The black bear in the spruce-fir forest. *Wildlife Monographs* 27. 57 pp.
- Kauffman, J. B., and W. C. Krueger. 1984. Livestock impacts on riparian ecosystems and streamside management implications—a review. *Journal of Range Management* 37:430-438.
- Kelleyhouse, D. G. 1980. Habitat utilization by black bears in northern California. *Bear Biology Association Conference Series* 3:221-228.
- Kiefling, J. W. 1978. Studies on the ecology of the Snake River cutthroat trout. *Wyoming Game and Fish Department Fisheries Technical Bulletin* 3. 192 pp.
- Knopf, F. L. 1985. Significance of riparian vegetation to breeding birds across an altitudinal cline. Pages 105-111 in R. R. Johnson, C. D. Ziebell, D. R. Patton, P. F. Ffolliott, and R. H. Hamre, technical coordinators. *Riparian ecosystems and their management: reconciling conflicting uses*. First North American Riparian Conference: 1985. U.S. Forest Service General Technical Report RM-120.
- Kochert, M. N., B. A. Millsap, and K. Steenhof. 1986. Effects of livestock grazing on raptors with emphasis on the southwestern U.S. Pages 325-333 in *Proceedings of the Southwest Raptor Management Symposium and Workshop*. National Wildlife Federation Scientific and Technical Series 11.
- Krausman, P. R., K. R. Rautenstrauch, and B. D. Leopold. 1985. Xeroriparian systems used by desert mule deer in Texas and Arizona. Pages 144-149 in *Riparian ecosystems and their management: reconciling conflicting uses*. U.S. Forest Service General Technical Report RM-120.
- Lee, L. C., T. A. Muir, and R. R. Johnson. 1987. Riparian ecosystems as essential habitat for raptors in the American West. Pages 15-26 in *Proceedings of the Western Raptor Management Symposium and Workshop*. National Wildlife Federation Scientific and Technical Series 12.
- Lingle, G. R. 1989. Winter raptor use of the Platte and North Platte river valleys in south central Nebraska. *Prairie Naturalist* 21:1-16.
- Marcus, M. D., M. K. Young, L. E. Noel, and B. A. Mullan. 1990. Salmonid-habitat relationships in the western United States: a review and indexed bibliography. U.S. Forest Service General Technical Report RM-188. 84 pp.
- Martin, P. R. 1977. The effect of altered stream flow on furbearing mammals of the Yellowstone River basin, Montana. *Montana Department of Natural Resource Conservation Technical Report* 6. 79 pp.
- Marzolf, G. R. 1978. The potential effects of clearing and snagging on stream ecosystems. U.S. Fish and Wildlife Service FWS/OBS-78/14. 32 pp.
- McArthur, J. V. 1989. Aquatic and terrestrial linkages: floodplain functions. Pages 107-116 in D. D. Hook and R. Lea, editors. *Proceedings of the symposium: the forested wetlands of the southern United States*. U.S. Forest Service General Technical Report SE-50.
- McGarigal, K., R. G. Anthony, and F. B. Isaacs. 1991. Interactions of humans and bald eagles on the Columbia River estuary. *Wildlife Monographs* 115. 47 pp.
- Medin, D. E., and W. P. Clary. 1989. Small mammal populations in a grazed and ungrazed riparian habitat in Nevada. U.S. Forest Service Research Paper INT-413. 6 pp.
- Melquist, W. E., and G. Hornocker. 1983. Ecology of river otters in west central Idaho. *Wildlife Monographs* 83. 60 pp.
- Melquist, W. E., J. S. Whitman, and M. G. Hornocker. 1981. Resource partitioning and coexistence of sympatric mink and river otter populations. Pages 187-220 in J. A. Chapman and D. Pursley, editors. *Proceedings of the Worldwide Furbearer Conference*, Frostburg, Md.
- Minshall, G. W., S. E. Jensen, and W. S. Platts. 1989. The ecology of stream and riparian habitats of the Great Basin region: a community profile. U.S. Fish and Wildlife Service Biological Report 85(7.24). 142 pp.
- Mitchell, J. L. 1961. Mink movements and populations on a Montana river. *Journal of Wildlife Management* 25:48-54.
- Mitsch, W. J., and J. G. Gosselink. 1986. *Wetlands*. Van Nostrand Reinhold Co., New York. 539 pp.
- Moulton, M. P., J. R. Choate, S. J. Bissel, and R. A. Nicholson. 1981. Associations of small mammals on the central high plains of eastern Colorado. *Southwestern Naturalist* 13:99-104.
- Munther, G. L. 1982. Beaver management in grazed riparian ecosystems. *Proceedings of the Wildlife-Livestock Relationships Symposium* 10:324-341.
- Odum, E. 1978. Ecological importance of the riparian zone. Pages 2-4 in *Strategies for protection and management of floodplain wetlands and other riparian ecosystems*. U.S. Forest Service General Technical Report WO-12.
- Ohmart, R. D., and B. W. Anderson. 1986. Riparian habitat. Pages 169-193 in A. Y. Cooperrider, R. J. Boyd, and H. R. Stuart, editors. *Inventory and monitoring of wildlife habitat*. U.S. Bureau of Land Management, Denver, Colo.
- Ohmart, D. R., W. O. Deason, and C. Burke. 1977. A riparian case history: the Colorado River. Pages 35-47 in *Importance, preservation and management of riparian habitat: a symposium*. U.S. Forest Service General Technical Report RM-43.

- Olson, T. E., and F. L. Knopf. 1988. Patterns of relative diversity within riparian small mammal communities, Platte River watershed, Colorado. Pages 379-386 in R. C. Szaro, K. E. Severson, and D. R. Patton, technical coordinators. Management of amphibians, reptiles, and small mammals in North America. U.S. Forest Service General Technical Report RM-166.
- Pederson, R. J., A. W. Adams, and J. Skovlin. 1979. Elk management in Blue Mountain habitats. Oregon Department of Fish and Wildlife, Portland. 27 pp.
- Platts, W. S., W. F. Megahan, and G. W. Minshall. 1983. Methods for evaluating stream, riparian, and biotic conditions. U.S. Forest Service General Technical Report INT-138. 70 pp.
- Platts, W. S., K. A. Gebhardt, and W. L. Jackson. 1985. The effects of storm events on basin-range riparian stream habitat. Pages 30-34 in R. Johnson, C. D. Ziebell, D. R. Patton, P. F. Ffolliott, and R. H. Hamre, technical coordinators. Riparian ecosystems and their management: reconciling conflicting uses. First North American Riparian Conference. U.S. Forest Service General Technical Report RM-120.
- Racey, G. D., and D. L. Euler. 1982. Small mammal and habitat response to shoreline cottage development in central Ontario. Canadian Journal of Zoology 60:865-880.
- Racey, G. D., and D. L. Euler. 1983. Changes in mink habitat and food selection as influenced by cottage development in central Ontario. Journal of Applied Ecology 20:387-402.
- Raedeke, K. J., R. D. Taber, and D. K. Paige. 1988. Ecology of large mammals in riparian systems of the Pacific Northwest forests. Pages 113-132 in K. J. Raedeke, editor. Streamside management: riparian wildlife and forestry interactions. University of Washington, Seattle.
- Reed, D. F. 1981. Conflicts with civilization. Pages 509-535 in O. C. Wallmo, editor. Mule and black-tailed deer of North America. University of Nebraska Press, Lincoln.
- Sampson, F. B., and F. L. Knopf. 1982. In search of a diversity ethic for wildlife management. Transactions of the North American Wildlife and Natural Resources Conference 47:421-431.
- Sanderson, G. C., E. D. Ables, R. D. Sparrow, J. R. Grieb, L. D. Harris, and A. N. Moen. 1979. Research needs in wildlife. Transactions of the North American Wildlife and Natural Resources Conference 44:166-175.
- Schroeder, R. L. 1987. Community models for wildlife impact assessment: a review of concepts and approaches. U.S. Fish and Wildlife Service Biological Report 87(2). 41 pp.
- Schroeder, R. L., T. T. Cable, and S. L. Haire. 1992. Wildlife species richness in shelterbelts: test of a habitat model. Wildlife Society Bulletin. In press.
- Sedell, J. R., and J. L. Froggatt. 1984. Importance of streamside forests to large rivers: the isolation of the Willamette River, Oregon, USA, from its floodplain by snagging and streamside forest removal. Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie 22:1828-1834.
- Sedell, J. R., and F. J. Swanson. 1984. Ecological characteristics of streams in old-growth forests of the Pacific Northwest. Pages 9-16 in W. R. Meehan, T. R. Merrell, Jr., and T. A. Hanley, editors. Fish and wildlife relationships in old-growth forests: proceedings of a symposium held in Juneau, Alaska 12-15 April 1982. American Institute of Fisheries Research Biologists.
- Simpson, J. C., and R. L. Wallace. 1982. Fishes of Idaho. University Press of Idaho, Moscow. 237 pp.
- Stauffer, D. F., and L. B. Best. 1980. Habitat selection by birds of riparian communities: evaluating effects of habitat alterations. Journal of Wildlife Management 44:1-15.
- Stevens, L. E., B. T. Brown, J. M. Simpson, and R. R. Johnson. 1977. The importance of riparian habitat to migrating birds. Pages 156-164 in R. R. Johnson and D. A. Jones, technical coordinators. Importance, preservation and management of riparian habitat. U.S. Forest Service General Technical Report RM-43.
- Stuber, R. J. 1985. Trout habitat, abundance, and fishing opportunities in fenced vs. unfenced riparian habitat along Sheep Creek, Colorado. Pages 310-314 in R. R. Johnson, D. D. Ziebell, D. R. Patton, P. F. Ffolliott, and R. H. Hamre, technical coordinators. Riparian ecosystems and their management: reconciling conflicting uses. Proceedings of the North American Riparian Conference. U.S. Forest Service General Technical Report RM-120.
- Szaro, R. C. 1980. Factors influencing bird populations in southwestern riparian forests. Pages 403-418 in Management of western forests and grasslands for nongame birds. U.S. Forest Service General Technical Report INT-86.
- Szaro, R. C., and J. N. Rinne. 1988. Ecosystem approach to management of southwestern riparian communities. Transactions of the North American Wildlife and Natural Resources Conference 53:502-511.
- U.S. Fish and Wildlife Service. 1990. Fish and Wildlife Coordination Act report for the Jackson Hole/Snake River flood protection/levee maintenance project, Teton County, Wyoming. U.S. Fish and Wildlife Service, Denver, Colo. 108 pp.

Vahle, J. R., N. L. Dodd, and S. Nagiller. 1986. Osprey. Pages 37-47 in *Proceedings of the southwest raptor management symposium and workshop*. National Wildlife Federation Scientific and Technical Series 11.

Warren, P. L., and C. R. Schwalbe. 1985. Herpetofauna in riparian habitats along the Colorado River in

Grand Canyon. Pages 347-354 in *Riparian ecosystems and their management: reconciling conflicting uses*. U.S. Forest Service General Technical Report RM-120.11.

Wilson, E. O. 1985. The biological diversity crisis. *BioScience* 35:700-706.

Appendix. Riparian-dependent vertebrates from the region of the Snake River near Jackson, Wyoming^a

Common name	Scientific name
Amphibians	
Tiger salamander	<i>Ambystoma tigrinum</i>
Boreal chorus frog	<i>Pseudacris triseriata maculata</i>
Northern leopard frog	<i>Rana pipiens</i>
Spotted frog	<i>Rana pretiosa</i>
Reptiles	
Wandering garter snake ^b	<i>Thamnophis elegans vagrans</i>
Valley garter snake ^b	<i>Thamnophis sirtalis fitchi</i>
Birds	
Great blue heron	<i>Ardea herodias</i>
Green-backed heron	<i>Butorides striatus</i>
Wood duck	<i>Aix sponsa</i>
Northern pintail	<i>Anas acuta</i>
American widgeon	<i>Anas americana</i>
Northern shoveler	<i>Anas clypeata</i>
Green-winged teal	<i>Anas crecca</i>
Cinnamon teal	<i>Anas cyanoptera</i>
Blue-winged teal	<i>Anas discors</i>
Mallard ^b	<i>Anas platyrhynchos</i>
Gadwall	<i>Anas strepera</i>
Lesser scaup	<i>Aythya affinis</i>
Redhead	<i>Aythya americana</i>
Ring-necked duck	<i>Aythya collaris</i>
Bufflehead	<i>Bucephala albeola</i>
Common goldeneye	<i>Bucephala clangula</i>
Barrow's goldeneye	<i>Bucephala islandica</i>
Trumpeter swan	<i>Cygnus buccinator</i>
Harlequin duck	<i>Histrionicus histrionicus</i>
Hooded merganser	<i>Lophodytes cucullatus</i>
Common merganser	<i>Mergus merganser</i>
Ruddy duck	<i>Oxyura jamaicensis</i>
Osprey	<i>Pandion haliaetus</i>
Broad-winged hawk	<i>Buteo platypterus</i>
Bald eagle	<i>Haliaeetus leucocephalus</i>
American coot ^b	<i>Fulica americana</i>
Spotted sandpiper ^b	<i>Actitis macularia</i>
Yellow-billed cuckoo	<i>Coccyzus americanus</i>
Black-billed cuckoo	<i>Coccyzus erythrophthalmus</i>

Appendix A. Continued.

Common name	Scientific name
Willow flycatcher	<i>Empidonax traillii</i>
Great crested flycatcher	<i>Myiarchus crinitus</i>
Bank swallow ^b	<i>Riparia riparia</i>
Northern rough-winged swallow ^b	<i>Stelgidopteryx serripennis</i>
Marsh wren	<i>Cistothorus palustris</i>
American dipper ^b	<i>Cinclus mexicanus</i>
Black-throated blue warbler	<i>Dendroica caerulescens</i>
Bay-breasted warbler	<i>Dendroica castanea</i>
Blackburnian warbler	<i>Dendroica fusca</i>
Palm warbler	<i>Dendroica palmarum</i>
Common yellowthroat	<i>Geothlypis trichas</i>
Yellow-breasted chat	<i>Icteria virens</i>
MacGillivray's warbler	<i>Oporornis tolmiei</i>
American redstart	<i>Setophaga ruticilla</i>
Wilson's warbler	<i>Wilsonia pusilla</i>
Summer tanager	<i>Piranga rubra</i>
Indigo bunting	<i>Passerina cyanea</i>
Lincoln's sparrow	<i>Melospiza lincolnii</i>
Song sparrow ^b	<i>Melospiza melodia</i>
Field sparrow	<i>Spizella pusilla</i>
White-crowned sparrow ^b	<i>Zonotrichia leucophrys</i>
American goldfinch ^b	<i>Carduelis tristis</i>
Masked shrew	<i>Sorex cinereus</i>
Mammals	
Northern water shrew	<i>Sorex palustris</i>
Vagrant shrew	<i>Sorex vagrans</i>
Mink	<i>Mustela vison</i>
River otter	<i>Lutra canadensis</i>
Moose	<i>Alces alces</i>
White-tailed deer	<i>Odocoileus virginianus</i>
Beaver	<i>Castor canadensis</i>
Montane vole ^b	<i>Microtus montanus</i>
Meadow vole ^b	<i>Microtus pennsylvanicus</i>
Water vole	<i>Microtus richardsoni</i>
Muskrat ^b	<i>Ondatra zibethicus</i>
Western jumping mouse	<i>Zapus princeps</i>

^a Adapted from Gerhart and Olson (1982) and U.S. Fish and Wildlife Service (1990).^b Species assumed to tolerate high levels of human disturbance.

A list of current *Resource Publications* follows.

171. Sand and Gravel Pits as Fish and Wildlife Habitat in the Southwest, by William J. Matter and R. William Mannan. 1988. 11 pp.
172. Satellite Telemetry: A New Tool for Wildlife Research and Management, by Steven G. Fancy, Larry F. Pank, David C. Douglass, Catherine H. Curby, Gerald W. Garner, Steven C. Amstrup, and Wayne L. Regelin. 1988. 54 pp.
173. Key to Acanthocephala Reported in Waterfowl, by Malcolm E. McDonald. 1988. 45 pp.
174. Obsolete English Names of North American Birds and Their Modern Equivalents, by Richard C. Banks. 1988. 37 pp.
175. Procedures for the Analysis of Band-recovery Data and User Instructions for Program MULT, by Michael J. Conroy, James E. Hines, and Byron K. Williams. 1989. 61 pp.
176. Sago Pondweed (*Potamogeton pectinatus* L.): A Literature Review, by Harold A. Kantrud. 1990. 89 pp.
177. Field Manual for the Investigation of Fish Kills, by Fred P. Meyer and Lee A. Barclay. 1990. 120 pp.
178. Section 404 and Alterations in the Platte River Basin of Colorado, by Douglas N. Gladwin, Mary E. Jennings, James E. Roelle, and Duane A. Asherin. 1990. 19 pp.
179. Hydrology of the Middle Rio Grande From Velarde to Elephant Butte Reservoir, New Mexico, by Thomas F. Bullard and Stephen G. Wells. 1991. 56 pp.
180. Waterfowl Production on the Woodworth Station in South-central North Dakota, 1965-1981, by Kenneth F. Higgins, Leo M. Kirsch, Albert T. Klett, and Harvey W. Miller. 1992. 79 pp.
181. Trends and Management of Wolf-Livestock Conflicts in Minnesota, by Steven H. Fritts, William J. Paul, L. David Mech, and David P. Scott. 1992. 27 pp.
182. Selection of Prey by Walleyes in the Ohio Waters of the Central Basin of Lake Erie, 1985-1987, by David Wolfert and Michael T. Burr. 1992. 14 pp.
183. Effects of the Lampricide 3-Trifluoromethyl- 4-Nitrophenol on the Pink Heelsplitter, by Terry D. Bills, Jeffrey J. Rach, Leif L. Marking, and George E. Howe. 1992. 7 pp.
184. Methods for Detoxifying the Lampricide 3-Trifluoromethyl-4-Nitrophenol in a Stream, by Philip A. Gilderhus, Terry D. Bills, and David A. Johnson. 1992. 5 pp.
185. Group Decision-making Techniques for Natural Resource Management Applications, by Beth A. K. Coughlan and Carl L. Armour. 1992. 55 pp.
186. DUCKDATA: A Bibliographic Data Base for North American Waterfowl (Anatidae) and Their Wetland Habitats, by Ken Reinecke and Don Delnicki. 1992. 8 pp.
187. Dusky Canada Goose: An Annotated Bibliography, by Bruce H. Campbell and John E. Cornely. 1992. 29 pp.
188. Human Disturbances of Waterfowl: An Annotated Bibliography, by Robert B. Dahlgren and Carl E. Korschgen. 1992. 62 pp.
189. Opportunities to Protect Instream Flows and Wetland Uses of Water in Nevada, by James L. Bingham and George A. Gould. 1992. 33 pp.

NOTE: Use of trade names does not imply U.S. Government endorsement of commercial products.

TAKE PRIDE

in America



U.S. DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE



As the nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our lands and water resources, protecting our fishes and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and attempts to assure that their development is in the best interests of all our people. The Department also has a major responsibility for Native American reservation communities and for people who live in island territories under U.S. administration.